

Package: Pade (via r-universe)

October 21, 2026

Type Package

Title Padé Approximant Coefficients

Version 1.1.0

Date 2026-03-25

Description Given a vector of Taylor series coefficients of sufficient length as input, the function returns the numerator and denominator coefficients for the Padé approximant of appropriate order (Baker, 1975) <ISBN:9780120748556>.

License GPL (>= 2) | BSD_2_clause + file LICENSE

Suggests covr, tinytest

URL <https://github.com/aadler/Pade>

BugReports <https://github.com/aadler/Pade/issues>

Encoding UTF-8

NeedsCompilation no

Repository <https://aadler.r-universe.dev>

Date/Publication 2026-03-25 05:52:28 UTC

RemoteUrl <https://github.com/aadler/pade>

RemoteRef HEAD

RemoteSha 1772767ab5a6840f819be2444036fa8fcecc789a

Contents

Pade-package	2
Pade	2
Index	5

Padé-package	<i>Padé Approximant Coefficients</i>
--------------	--------------------------------------

Description

Given a vector of Taylor series coefficients of sufficient length as input, the function returns the numerator and denominator coefficients for the Padé approximant of appropriate order (Baker, 1975) <ISBN:9780120748556>.

Details

The DESCRIPTION file: This package was not yet installed at build time.

Index: This package was not yet installed at build time.

Author(s)

Avraham Adler [aut, cph, cre] (ORCID: <<https://orcid.org/0000-0002-3039-0703>>)
Maintainer: Avraham Adler <Avraham.Adler@gmail.com>

Padé	<i>Padé Approximant Coefficients</i>
------	--------------------------------------

Description

Given Taylor series coefficients a_n from $n = 0$ up to $n = T$, the function will calculate the Padé $[L/M]$ approximant coefficients so long as $L + M \leq T$.

Usage

```
Padé(L, M, A, method = c("qr", "solve"), tol = 1e-7)
```

Arguments

- | | |
|--------|--|
| L | numeric ; scalar order of Padé numerator |
| M | numeric ; scalar order of Padé denominator |
| A | numeric ; vector of Taylor series coefficients, starting at x^0 |
| method | character ; selects linear-equation solver algorithm. qr is more robust to ill-conditioned matrices. solve is faster. |
| tol | numeric ; the tolerance passed to qr.solve. Defaults to 1e-7, the default value in qr.solve. |

Details

As the Taylor series expansion is the “best” polynomial approximation to a function, the Padé approximants are the “best” rational function approximations to the original function. The Padé approximant often has a wider radius of convergence than the corresponding Taylor series, and can even converge where the Taylor series does not. This makes it very suitable for computer-based numerical analysis.

The $[L/M]$ Padé approximant to a Taylor series $A(x)$ is the quotient

$$\frac{P_L(x)}{Q_M(x)}$$

where $P_L(x)$ is of order L and $Q_M(x)$ is of order M . In this case:

$$A(x) - \frac{P_L(x)}{Q_M(x)} = \mathcal{O}(x^{L+M+1})$$

When q_0 is defined to be 1, there is a unique solution to the system of linear equations which can be used to calculate the coefficients.

The function accepts a vector A of length $T + 1$, composed of the a_n of the truncated Taylor series

$$A(x) = \sum_{j=0}^T a_j x^j$$

and returns a list of two elements, Px and Qx , the Padé numerator and denominator coefficients respectively, as long as $L + M \leq T$.

Value

Pade returns a list with two entries:

Px	Coefficients of the numerator polynomial starting at x^0 .
Qx	Coefficients of the denominator polynomial starting at x^0 .

Author(s)

Avraham Adler <Avraham.Adler@gmail.com>

References

Baker, George Allen (1975) *Essentials of Padé Approximants* Academic Press. ISBN 978-0-120-74855-6

See Also

This package provides similar functionality to the `pade` function in the **pracma** package. However, it does not allow computation of coefficients beyond the supplied Taylor coefficients and it expects its input and provides its output in ascending—instead of descending—order.

See the **minimaxApprox** package for polynomial and rational minimax approximations to functions.

Examples

```
A <- 1 / factorial(0:10) ## Taylor sequence for e^x up to x^{10} around x_0 = 0
Z <- Pade(5, 5, A)
print(Z)                    ## Padé approximant of order [5 / 5]
x <- -.01                   ## Test value
Actual <- exp(x)            ## Proper value
print(Actual, digits = 16)
Estimate <- sum(Z[[1L]] * x ^ (seq_along(Z[[1L]]) - 1)) /
  sum(Z[[2L]] * x ^ (seq_along(Z[[2L]]) - 1))
print(Estimate, digits = 16) ## Approximant value
all.equal(Actual, Estimate)
```

Index

- * **NumericalMathematics**

- Pade, [2](#)

- Pade-package, [2](#)

- * **algebra**

- Pade, [2](#)

- * **math**

- Pade, [2](#)

- * **package**

- Pade-package, [2](#)

Pade, [2](#)

Pade-package, [2](#)